

GROUP 4

Active Ingredients

Fluroxypyr 90 g/L and 2,4-D ester 360 g/L as an EC

Packaging

Case: 2 x 8.9 L jug Bulk: 106.8 L drums Tote: 427.2 L

Application Rates & Acres Treated

Rate: 445 ml/ac Acres Treated: 40 ac/case 240 ac/drum 960 ac/tote

Water Volume

Ground: 40 L/ac

(10 US gal/ac)

Aerial: Do not apply.

Rainfastness

1 hour

KEY BENEFITS

- Control of Group 2-resistant biotypes of kochia
- Suppression of Group 2-resistant biotypes of chickweed
- Convenient and trusted formulation
- Easily tankmixes with the most popular grassy weed control herbicides
- Available in convenient bulk options

REGISTERED CROPS

Barley

Wheat (durum, spring)

WEEDS CONTROLLED

Suppression At labelled rate of 445 ml/ac Common Chickweed² Bluebur Mustards (except Green Burdock Tansy, Dog & Grey Tansy) (up to 8 cm) Canola (Volunteer) Plantain Redroot Pigweed Prickly Lettuce • Sow Thistle (Perennial)¹ Cleavers (1-8 whorls) Clovers (Sweet) Ragweed Cocklebur Round-leaved Mallow Flixweed (1- to 6-leaf) Field Horsetail¹ Shepherd's Purse Goat's-beard Stork's Bill (1- to 8-leaf) Stinkweed Hemp-nettle (2- to 6-leaf stage) Sunflower (Annual) Hoary Cress¹ Vetch Kochia² Wild Radish Lamb's Quarters • Volunteer Flax (1-12 cm) Additional weeds controlled/suppressed with addition of 81 ml/ac (2 oz/ac) of 2,4-D ESTER 700: Blue Lettuce¹ Lady's Thumb Common Chickweed² • Dandelion (Spring Leafy Spurge¹ (up to 8 cm) Mustard (Dog, Tansy) Canada Thistle¹ Rosettes) Docks Oak-leaved Goosefoot Sow Thistle Field Bindweed¹ Redroot Pigweed (Perennial¹, Annual) • Field Peppergrass • Russian Thistle Gumweed Smartweed Hairy Galinsoga Tartary Buckwheat

Hedge Bindweed
 Top growth control only

² Including biotypes resistant to Group 2 herbicides that inhibit the ALS enzyme.



RUSH 24 ALL IN™

HOW IT WORKS

Group 4 herbicides disrupt normal plant growth, resulting in twisting and cupping of leaves, epinasty and death of susceptible plants in 2–10 days.

REGISTERED AND SUPPORTED TANK MIXES

Wheat only:

- Everest® 3.0
 Simplicity™ GoDRI™ 1
 Varro®
- LADDER ALL IN®
 Traxos®

Wheat and barley:

Assert® 300 SC
 BISON® 400 L
 Puma® Advance

MIXING INSTRUCTIONS

- 1. Fill the clean spray tank ½ full of clean water.
- 2. Add the required amount of RUSH 24 ALL IN™ and agitate thoroughly, followed by the required amount of additional 2,4-D ESTER 700 herbicide, if required.
- **3.** Continue agitation and add the required amount of the tank-mix partner.
- 4. Complete filling the tank to the desired level with water.

CROP ROTATIONS

The following crops may be grown one (1) year after application:

- BarleyCanolaMustard
- FlaxForage grassField peas

PRE-HARVEST INTERVAL

Barley, wheat (durum, spring): 60 days

STORAGE

Do not freeze.

GRAZING RESTRICTIONS

- Do not permit lactating dairy animals to graze fields within seven (7) days after application.
- Do not harvest forage or cut hay within 30 days after application.
- Withdraw meat animals from treated fields at least three (3) days before slaughter.



Rve

Wheat



Quick Tips

RUSH 24 ALL IN™ activity is influenced by weather conditions. Optimum activity requires active crop and weed growth. The temperature range for optimum activity is 12°C to 24°C. Reduced activity will occur when temperatures are below 8°C or above 27°C. Frost before application (3 days) or shortly after (3 days) may reduce weed control and crop tolerance. Weed control may be reduced during stress conditions, for example drought, heat or cold stress, or if weeds have initiated flowering, or if heavy infestations exist.



Always read and follow label directions.
Toll-free:1.855.264.6262 | Website: ADAMA.COM

¹ Additional 2,4-D ester is not recommended when mixing RUSH 24 ALL IN™ and Simplicity™.