



ADAMA

Custodia® Forte

Fungicide

Product overview

Custodia® Forte fungicide provides protectant and systemic activity against the key diseases in almonds, grapes and macadamias. Custodia Forte is a concentrated suspension concentrate formulation that combines two different modes of action (Group 3 + 11) for robust disease control and resistance management.

Key features

- Preventative control or suppression of key diseases protecting growers yield potential and quality
- Two modes of action for improved disease control and resistance management
- Excellent protectant and systemic activity, suitable to use in IPM programs
- Suspension concentrate formulation with low odour
- Highly compatible formulation for tank mixing

Application rate conversion

Custodia Forte is a concentrated formulation containing 85% more active ingredient than Custodia, reducing on-farm packaging and waste.

Custodia (200 g/L tebuconazole + 120 g/L azoxystrobin)	Custodia Forte (370 g/L tebuconazole + 222 g/L azoxystrobin)
Dilute rate (mL/100 L water)	
60	32
100	54

Product information

Active ingredient:
Azoxystrobin 222 g/L +
Tebuconazole 370 g/L

Mode of action:
Group 3 + 11

Formulation:
Suspension Concentrate

Registered crops:
Almonds, macadamias, grapes

Target diseases:
Brown rot/Blossom blight,
Shot-hole, Rust, Hull rot,
Powdery mildew, Downy mildew,
Botrytis bunch rot, Husk spot

Re-entry period:
Once the spray has dried

Custodia® Forte

Crop	Target disease	Application rate	Timing	Spray frequency
Almonds	Brown Rot/Blossom blight and Shot-hole	32 mL/100 L or 540 mL/ha	From early (10%) bloom before early hull split	Apply in rotation with fungicides from alternative modes of action on a 7 to 14-day interval targeting key growth stages including full bloom, petal fall and shuck fall After shuck fall, apply Cavalry® Weatherguard for Shot-hole control on a 14-day spray interval Do not apply consecutive applications and no more than three sprays per season
	Rust		Commence spray program from late flowering and/or when first new leaves emerge	Apply up to two applications per season in rotation with fungicides from a different mode of action group on a 14-day spray interval
	Hull rot (suppression only)		Early hull split	Apply a single application. If a second fungicide application is required, rotate with an alternative registered fungicide after a 10 to 14-day interval
Grapes	Powdery Mildew	35 mL/100 L	Before E-L 19 (before flowering) and under low disease pressure up to E-L 29	Maximum two applications per season. Always apply as a preventative spray and never as a curative spray
	Downy Mildew	54 mL/100 L or 540 mL/ha	From after E-L 19 (start of flowering) and under high disease pressure up to E-L 29	
	Botrytis Bunch Rot (low disease pressure only)		Seasonal preventative spray program commencing when shoots are 20 cm long, up to E-L 29	Apply in rotation with fungicides from other modes of action on a 7 to 21-day spray interval. Under high disease pressure and during flowering, apply in rotation with Axiom® Plus
			Use no later than E-L 29	If three or fewer Botrytis sprays are applied in a season, use no more than one Custodia Forte spray during the season If four or more Botrytis sprays are to be applied in a season, up to two Custodia Forte sprays can be applied
Macadamia	Husk spot	32 mL/100 L*	From match head timing	Two applications per season on a 14 to 28-day spray interval dependent on disease pressure, with fungicides from other modes of action groups.

Applications of Custodia Forte with Hasten as per label guidelines can improve husk spot control under conditions of high disease pressure. However, we advise growers to evaluate this mix in a small area before commercial use

Harvest Withholding Period

Almonds: Not required when used as directed

Grapevines: Do not harvest for 4 weeks after application

Macadamias: Do not harvest for 15 days after application



Scan here for more information

Refer to label for specific application recommendations.
™Trademarks or ®Registered trademarks of an ADAMA Agricultural Solutions Ltd Company.
 *Registered trademarks. Please note: This information is not intended to replace the product label.
 Always read the complete product label appearing on the container before opening or using products. ADA22186



ADAMA.COM

1800 4 ADAMA

