




ADAMA

TOPNOTCH™

A new broad spectrum, multi-mode of action option for control of dangerous diseases in dry beans and soybeans as well as cereal leaf diseases.



GROUP 3 & 11

Active Ingredient

Azoxystrobin 143 g/L and Propiconazole 124 g/L = SC

Packaging

Case: 2 x 8.6 L jugs

Application Rates & Acres Treated

Rate: 210 – 620 ml/ac

Acres Treated: 14–40 ac/jug

Water volume

Ground: 40 L/ac (10 US gal/ac)

Aerial: 20 L/ac (5 US gal/ac)

Rainfastness

Avoid applying when heavy rainfall is in the forecast.

KEY BENEFITS

- Multiple modes of action for resistance management
- Curative and preventative systemic action
- Can be used on multiple crops, with exceptional crop safety
- Registered for both ground and aerial spraying
- Excellent crown rust control in oats

KEY DISEASES CONTROLLED

- Anthracnose
- Ascochyta blight
- Barley leaf rust
- Mycosphaerella blight
- Net and spot blotches
- Powdery mildew
- Scald
- Septoria spot
- Stripe rust
- Tan spot
- Wheat leaf rust
- White mould¹

HOW IT WORKS

Used as both a curative and preventative fungicide, TOPNOTCH™ has broad-spectrum, systemic and contact activity.

APPLICATION TIMING AND CROP STAGING

Crop	Diseases	Application Timing	Rate
Barley	Barley net blotch, Barley scald, Septoria leaf spot, Stripe rust, Barley leaf rust, Tan spot	Apply once between stem elongation and half-head emergence (Growth stage 29–55).	210 ml/ac
Oats	Barley net blotch, Crown rust, Septoria leaf spot	Apply once between stem elongation and half-head emergence (Growth stage 29–55).	
Rye	Septoria leaf spot, Barley scald, Tan spot	Apply once between stem elongation and half-head emergence (Growth stage 29–55).	
Triticale	Septoria leaf spot, Tan spot		
Wheat	Septoria leaf spot, Tan spot, Stripe rust, Wheat leaf rust		




TOPNOTCH™


APPLICATION TIMING AND CROP STAGING (CONT'D)

Crop	Diseases	Application Timing	Rate
Beans, Field peas, Lentils, Soybeans	Mycosphaerella blight, Anthracnose	Make the first application at the first sign of disease. Apply the high rate only under conditions of high disease pressures. A second application 14 days later may be needed if conditions persist. Good spray coverage and canopy penetration are important for best results.	310 ml/ac – 620 ml/ac
	Powdery mildew, White mould (suppression only)		310 ml/ac

PEA GROWTH STAGES




SOYBEAN GROWTH STAGES




TOPNOTCH™

LENTILS GROWTH STAGES



CEREAL GROWTH STAGES



Timing based on using the product alone. Please refer to tank-mix partners label for crop timing.

Always read and follow label directions.



TOPNOTCH™

SUPPORTED TANK MIXES

Insecticides:

- SILENCER® 120 EC
- Coragen®
- Voliam XPress®
- Decis®

MIXING INSTRUCTIONS

1. Fill spray tank ½ – ¾ full with water.
2. With agitator running, add required amount of TOPNOTCH™ and continue agitating while adding remainder of the water.
3. Begin application after TOPNOTCH™ is completely dispersed into the mix water, and maintain agitation during spraying operation.

CROP ROTATIONS

Do not plant any other crop intended for food, grazing or any component of animal feed or bedding within 105 days of application.

PRE-HARVEST INTERVALS

Beans, Field peas, Lentils, Soybeans: 30 days
Cereals: 45 days

GRAZING RESTRICTIONS

No restrictions.

STORAGE

Do not freeze.

Quick Tips

Good spray coverage and canopy penetration are important to achieve the best results.



ADAMA

Always read and follow label directions.
1.855.264.6262 ADAMA.COM

®/™ BUMPER, CUSTODIA, ORIUS and SILENCER are registered trademarks, and TOPNOTCH is a trademark of ADAMA Agricultural Solutions Canada Ltd. All other products are trademarks of their respective companies.
© 2022 ADAMA Agricultural Solutions Canada Ltd.