



# RUSH 24 ALL IN™

Controls a wide spectrum of broadleaf weeds, like cleavers, with excellent grassy weed herbicide compatibility for wheat and barley



## GROUP 4

### Active Ingredients

Fluroxypyr 90 g/L and 2,4-D ester 360 g/L as an EC

### Packaging

Case: 2 x 8.9 L jugs

### Application Rates & Acres Treated

Rate: 445 ml/ac

Acres Treated:  
20 ac/jug

### Water Volume

Ground: 10 US gal/ac  
(40 L/ac)

Aerial: Do not apply.

### Rainfastness

1 hour

### KEY BENEFITS

- Control of Group 2-resistant biotypes of kochia
- Suppression of Group 2-resistant biotypes of chickweed
- Convenient and trusted formulation
- Easily tankmixes with the most popular grassy weed control herbicides
- Available in convenient bulk options

### REGISTERED CROPS

- Barley
- Wheat (spring)

### WEEDS CONTROLLED

Control		Suppression
At labelled rate of 445 ml/ac		
<ul style="list-style-type: none"> <li>• Bluebur</li> <li>• Burdock</li> <li>• Canola (volunteer)</li> <li>• Cleavers (1-8 whorls)</li> <li>• Clovers (sweet)</li> <li>• Cocklebur</li> <li>• Flixweed</li> <li>• Field horsetail<sup>1</sup></li> <li>• Goat's-beard</li> <li>• Hemp-nettle (2- to 6-leaf stage)</li> <li>• Hoary cress<sup>1</sup></li> <li>• Kochia<sup>2</sup></li> <li>• Lamb's quarters</li> </ul>	<ul style="list-style-type: none"> <li>• Mustards (except green tansy, dog &amp; grey tansy)</li> <li>• Plantain</li> <li>• Prickly lettuce</li> <li>• Ragweed</li> <li>• Round-leaved mallow (1- to 6-leaf)</li> <li>• Shepherd's purse</li> <li>• Stork's bill (1- to 8-leaf)</li> <li>• Stinkweed</li> <li>• Sunflower (annual)</li> <li>• Vetch</li> <li>• Wild radish</li> <li>• Volunteer flax (1-12 cm)</li> </ul>	<ul style="list-style-type: none"> <li>• Common chickweed<sup>2</sup> (up to 8 cm)</li> <li>• Redroot pigweed</li> <li>• Sow thistle (perennial)<sup>1</sup></li> </ul>
Additional weeds controlled/suppressed with addition of 81 ml/ac (2 oz/ac) of 2,4-D ESTER 700:		
<ul style="list-style-type: none"> <li>• Blue lettuce<sup>1</sup></li> <li>• Dandelion (spring rosettes)</li> <li>• Docks</li> <li>• Field bindweed<sup>1</sup></li> <li>• Field peppergrass</li> <li>• Gumweed</li> <li>• Hairy galinsoga</li> <li>• Hedge bindweed</li> </ul>	<ul style="list-style-type: none"> <li>• Lady's thumb</li> <li>• Leafy spurge<sup>1</sup></li> <li>• Mustard (dog, tansy)</li> <li>• Oak-leaved goosefoot</li> <li>• Redroot pigweed</li> <li>• Russian thistle</li> <li>• Smartweed</li> <li>• Tartary buckwheat</li> </ul>	<ul style="list-style-type: none"> <li>• Common chickweed<sup>2</sup> (up to 8 cm)</li> <li>• Canada thistle<sup>1</sup></li> <li>• Sow thistle (perennial<sup>1</sup>, annual)</li> </ul>

<sup>1</sup>Top growth control only

<sup>2</sup>Including biotypes resistant to Group 2 herbicides that inhibit the ALS enzyme.



# RUSH 24 ALL IN™

## HOW IT WORKS

Group 4 herbicides disrupt normal plant growth, resulting in twisting and cupping of leaves, epinasty and death of susceptible plants in 2–10 days.

## REGISTERED AND SUPPORTED TANK MIXES

### Wheat only:

- Simplicity™ GoDRI™<sup>1</sup>
- Traxos®

### Wheat and barley:

- BISON® 400 L
- Puma® Advance

<sup>1</sup> Additional 2,4-D ester is not recommended when mixing RUSH 24 ALL IN™ and Simplicity™.

## MIXING INSTRUCTIONS

1. Fill the clean spray tank ½ full of clean water.
2. Add the required amount of RUSH 24 ALL IN™ and agitate thoroughly, followed by the required amount of additional 2,4-D ESTER 700 herbicide, if required.
3. Continue agitation and add the required amount of the tank-mix partner.
4. Complete filling the tank to the desired level with water.

## CROP ROTATIONS

The following crops may be grown one (1) year after application:

- Barley
- Canola
- Forage grass
- Oats
- Field peas
- Rye
- Wheat

## PRE-HARVEST INTERVAL

Barley, wheat (spring): 60 days

## STORAGE

Do not freeze.

## GRAZING RESTRICTIONS

- Do not permit lactating dairy animals to graze fields within seven (7) days after application.
- Do not harvest forage or cut hay within 30 days after application.
- Withdraw meat animals from treated fields at least three (3) days before slaughter.

### QUICK TIPS:

RUSH 24 ALL IN™ activity is influenced by weather conditions. Optimum activity requires active crop and weed growth. The temperature range for optimum activity is 12°C to 24°C. Reduced activity will occur when temperatures are below 8°C or above 27°C. Frost before application (3 days) or shortly after (3 days) may reduce weed control and crop tolerance. Weed control may be reduced during stress conditions, for example drought, heat or cold stress, or if weeds have initiated flowering, or if heavy infestations exist.

## WANT TO KNOW MORE?

Click on the QR code to access all the product details including the label and SDS documents



Always read and follow label directions.  
1.855.264.6262 ADAMA.COM

®/™BISON is registered trademarks and RUSH ALL IN is a trademark of ADAMA Agricultural Solutions Canada Ltd.  
All other products are trademarks of their respective companies.  
© 2024 ADAMA Agricultural Solutions Canada Ltd.

1/2024