



ADAMA

# VANTANA™

A protective, broad-spectrum fungicide used in a variety of fruit and vegetable crops. This Group 29 fungicide is a very important resistance management tool with activity on the hardest-to-control fungal diseases.



## Active Ingredient

Fluazinam 500 g/L = SC

## Packaging

Case: 2 x 10 L jugs

## Application Rates & Acres Treated

Rate: 162 – 906 ml/ac

Acres Treated: 27 – 80 ac/jug

## Water Volume

**Ground:** Varies with the crop. Refer to the label for more details.

**Aerial:** Minimum of 18.2 L/ac

## Rainfastness

Avoid application when heavy rain is forecast.

## GROUP 29

## KEY BENEFITS

- Resistance Management Tool
- Can be used on multiple crops
- Broad spectrum disease control
- Multiple rates for precise control

## KEY REGISTERED CROPS

- Apples
- Blueberries (highbush, lowbush)
- Bulb onion
- Currant
- Dry-shelled beans
- Elderberry
- Edible-podded legume vegetables (except peas)
- Ginseng
- Gooseberry
- Huckleberry
- Head and Stem Brassica (Crop Subgroup 5A)
- Leafy Brassica Greens (Crop Subgroup 5B)
- Potatoes
- Soybeans

For a full list of registered crops, please refer to the label.

## HOW IT WORKS

VANTANA™ works by inhibiting fungal adenosine triphosphate (ATP) production in the mitochondria (impairing energy production).

Active on a wide range of diseases, VANTANA™ is a protective fungicide that inhibits the germination of fungal spores.

## MIXING INSTRUCTIONS

1. Add ½ of the required amount of water to the spray or mixing tank and start agitation.
2. Add the required quantity of VANTANA™ to the water and complete filling with water to the required total volume.
3. Maintain agitation throughout mixing and spraying.

NOTE: If tank-mixing with an insecticide, please reference the label of the partner for specific mixing order or follow W.A.M.L.E.G.S or W.A.L.E.S. for proper mixing protocol.



# VANTANA™

## KEY DISEASES AND RATES BY CROP

NOTE: For application timing by crop, please refer to the label.

Crop	Diseases	Rate
Apples	Cedar apple rust	304–405 ml/ac (750–1,000 ml/ha)
	<i>Suppression:</i> Black rot, Brooks spot, Quince rust	405 ml/ac (1,000 ml/ha)
Blueberries (lowbush)	Valdensinia leaf spot	162–324 ml/ac (400–800 ml/ha)
Blueberries (highbush, lowbush)	<i>Suppression:</i> Mummyberry, Phomopsis fruit rots and fruit anthracnose	907 ml/ac (2,240 ml/ha)
	<i>Suppression:</i> Fruit anthracnose	
Bulb Onion	Purple blotch, Botrytis leaf blight	469 ml/ac (116 ml/ha) (584 g)
Dry-shelled beans	White mould	243–405 ml/ac (600–1,000)
	Anthracnose	
Edible-podded legume vegetables (except pea)	White mould	
Ginseng	Rhizoctonia root rot	486 ml/ac (1,200 ml/ha)
	Alternaria blight, Botrytis blight	
Potatoes	Late blight	162 ml/ac (400 ml/ha)
	White mould	162–242 ml/ac (400–600 ml/ha)
Soybeans	White mould	356–473 ml/ac (880–1,170 ml/ha)
	<i>Suppression:</i> White mould	178 ml/ac (440 ml/ha)
Leafy Brassica Greens (Crop Subgroup 5B) <b>Pre-transplant treatment</b>	Clubroot	1.17 L in 202 L of water per acre (2.9 L in 500 L of water per ha)
Head and Stem Brassica (Crop Subgroup 5A), Leafy Brassica Greens (Crop Subgroup 5B) <b>Transplant treatment</b>		Mix 50 ml with water to make a 100 L solution.

## CROP ROTATIONS

Areas treated with VANTANA™ may be replanted with potatoes and dry shelled beans as soon as practical after the last application. Other root crops and leafy vegetables can be planted 30 days after the last application. All other crops can be planted 70 days after the last application.

## GRAZING RESTRICTIONS

Do not feed treated foliage to livestock.

## PRE-HARVEST INTERVALS

Bulb onion: 7 days

Edible-podded legume vegetables (except peas),

Potatoes: 14 days

Apples: 28 days

Blueberries (highbush, lowbush), Currant, Dry-shelled beans, Elderberry, Ginseng, Gooseberry, Huckleberry  
Leafy Brassica Greens (Crop Subgroup 5B): 30 days

Head and Stem Brassica (Crop Subgroup 5A): 65 days

Soybeans: DO NOT apply after growth stage R3

## STORAGE


Do not freeze.

### QUICK TIP

For optimal efficacy VANTANA™ requires good coverage, best achieved with higher water volumes. Actual water volumes will vary with the crop, please refer to the label for more specific details.

## WANT TO KNOW MORE?

Click on the QR code to access all the product details including the label and SDS documents



ADAMA

Always read and follow label directions.  
1.855.264.6262 ADAMA.COM

®/™ VANTANA is a trademark of ADAMA Agricultural Solutions Canada Ltd.  
All other products are trademarks of their respective companies.  
© 2024 ADAMA Agricultural Solutions Canada Ltd.  
02/2024