

Paranex™

Paraquat Dichloride 24% SL

- ▶ **Paranex** belongs to bipyridylium group.
- ▶ **Paranex** is a non-selective, fast acting, contact herbicide, absorbed by the foliage.
- ▶ **Paranex** acts in the presence of light to desiccate the green parts of weed plant with which it comes into contact. It does not affect the brown mature stem and bark of tree. The product is very fast in action and controls weeds within 48 hours.



Available Pack sizes:
250ml, 500ml, 1ltr, 5ltr, 20ltr

Antidote: No specific antidote is known. Treat symptomatically. 500 ml of 30 % suspension of sterilized bentonite or fullers earth may be given orally



Recommendation for use:

Crop	Weeds	Dosage	
		Lt/Hectare	Lt/Acre
Tea (Post-emergence directed inter row application at 2-3 leaf stage of weeds)	<i>Imperata cylindrica</i> , <i>Setaria sp.</i> , <i>Commelina benghalensis</i> , <i>Boerhavia hispida</i> , <i>Paspalum conjugatum</i>	0.8-5	0.32-2
Potato (Post-emergence overall / inter-row application at 5-10 % emergence)	<i>Chenopodium sp.</i> , <i>Angallis arvensis</i> , <i>Triathema monogyna</i> , <i>Cyperus rotundus</i> , <i>Fumaria parviflora</i>	2	0.8

Paranex™

Paraquat Dichloride 24% SL

Recommendation for use:

Crop	Weeds	Dosage	
		Lt/Hectare	Lt/Acre
Cotton (Post-emergence directed inter row application at 2-3 leaf stage of weeds)	<i>Digera arvensis</i> , <i>Cyperus iria</i> , <i>Trianthema portulacastrum</i>	1.25-2.0	0.5-0.8
Rubber (Post-emergence directed inter row application at 2-3 leaf stage of weeds)	<i>Digitaria sp.</i> , <i>Eragrostis sp.</i> , <i>Fimbristylis sp.</i> ,	1.5-2.5	0.60-1
Rice [pre-plant (minimum tillage) before sowing / transplanting for controlling standing weeds]	<i>Ageratum conyzoides</i> , <i>Commelia benghalensis</i> , <i>Echinochloa crusgalli</i> , <i>Panicum repens</i> , <i>Cyperus iria</i> , <i>Brachiaria mutica</i> , <i>Marsilea quadriifoliata</i> ,	1.25-3.5	0.5-1.4
Wheat [pre-plant (minimum tillage) before sowing]	Grassy & Broad leaf weeds	5	2
Maize [pre-plant (minimum tillage) before sowing]	<i>Cyperus rotundus</i> , <i>Commelina benghalensis</i> , <i>Trianthema monogyna</i> , <i>Amaranthus sp.</i> , <i>Echinochloa sp</i>	0.8-2.0	0.32-0.8
Grape (Post-emergence directed inter row application at 2-3 leaf stage of weeds)	<i>Cyperus rotundus</i> , <i>Cynodon dactylon</i> <i>Convolvulus sp.</i> , <i>Portulaca sp.</i> , <i>Tridax sp.</i>	2	0.8
Apple (Post-emergence directed inter row application at 2-3 leaf stage of weeds)	<i>Rosa moschata</i> , <i>Rosa eglanteria</i> , <i>Rubus ellipticus</i>	3.25	1.3

Paranex™

Paraquat Dichloride 24% SL

Recommendation for use:

Crop	Weeds	Dosage	
		Lt/Hectare	Lt/Acre
Coffee	<i>Digitaria marginata, paspalum Conjugatum, Ageratum, Conyzides, Borreria hispida, Euphorbia hirta, Commelina benghalensis, Eleusine indica</i>	1	0.4
Sugarcane	<i>Echinochloa crusgalli, Digiteria sanguinalis, Eleusine indica, Physalis minima, Portulaca oleracea.</i>	2	0.8
Sunflower	<i>Cyperus iria, Eleusine indica, Eragrostis sp., Chenopoduim album, Trianthema monogyna</i>	1.6	0.64
Aquatic weed control	<i>Eichhornia crassipes</i>	4.25	1.7
	<i>Hydrilla</i>	4.25	1.7
Water ways Canals, Ponds	<i>Typha latifolia</i>	4.25-8.5	1.7-3.4