



# VICTOR GEL

Reg no. L8594 Act/Wet 36 of/van 1947

A non-volatile, non-repellent gel bait insecticide for the control of German cockroaches in domestic and food storage premises, as well as in public hygiene. 'n Nie-vlugtige, nie-afwerende jel insekdoder vir die beheer van Duitse kakkerlakke in huishoudelike persele en voedselopbergings-fasiliteite asook in publieke higiënesituasies.

**IRAC INSECTICIDE GROUP CODE**                      **4A**                      **IRAC INSEKDODER GROEPKODE**

## ACTIVE INGREDIENT/AKTIEWE BESTANDDEEL

Imidacloprid (chloronicotinyl) ..... 21.5 g/kg ..... Imidaklopried (chloronikotiniel)

## NET MASS/NETTO MASSA

..... g

## REGISTRATION HOLDER/REGISTRASIEHOUER

ADAMA South Africa (Pty.) Ltd.

Reg. no. 1992/001741/07

21 Viben Street

7560 Brackenfell

T: +27 21 982 1460 F: +27 21 982 5810

infocpt@adama.com

**VICTOR GEL is the trademark of a company of the ADAMA GROUP.**  
**VICTOR GEL is die handelsmerk van 'n maatskappy van die ADAMA GROEP.**

**ONLY FOR SALE TO AND USE BY REGISTERED PEST CONTROL OPERATORS**  
**SLEGS VIR VERKOOP AAN EN GEBRUIK DEUR GEREGISTREERDE PLAAGBEHEEROPERATEURS**

## CONTACT IN EMERGENCY/KONTAK IN NOOD

Griffon Poison Information Centre: +27 82 446 8946

Batch number .....  
Date of Manufacture .....

Lotnommer .....  
Datum van Vervaardiging .....

UN No.:



## **WARNINGS**

- Handle with care.
- Harmful when swallowed.
- Harmful to fish and other animals.
- Store in a cool place away from food and feedstuffs.
- Keep out of reach of children, uninformed persons and animals.

## **Precautions**

- Avoid contact by mouth or with skin.
- Wear rubber gloves during application.
- Wash with soap and water immediately after use or after accidental skin contact.
- Do not eat, drink or smoke whilst mixing, applying or before washing hands and face.
- Apply only in area where it will be completely inaccessible to children and pets.
- Do not contaminate dams, drains, ponds, streams and other waterways with the product or with used containers.
- Do not apply to surfaces on which food or feed is stored, prepared or eaten.
- Protect food preparing equipment and eating utensils from contamination during application.
- Destroy the empty container and do not re-use for any other purpose.

## **DIRECTIONS FOR USE**

### **Use only as indicated.**

Suitable for use in commercial and industrial premises, warehouses, food handling and processing establishments, market places, restaurants, hotels, hospitals, barracks, ships, offices, store rooms, schools and homes, etc.

**VICTOR GEL** is a cockroach insecticide with delayed action; therefore the results will not be visible immediately. A noticeable effect on the cockroach population may only be observed within a week. The area to undergo treatment should be examined to identify the centres of the cockroach infestation. **VICTOR GEL** can then be applied by spot-treatment in places in and around cockroach harborage and where cockroaches have been observed or where they hide. The gel should be applied in dark areas and well out of sight.

**VICTOR GEL** is dispensed from a tube using a purpose-built gun, which is operated as follows:

Insert the cartridge/tube into the gun frame. Press the trigger until the plunger contracts the piston of the cartridge but do not apply pressure. Remove the outer nozzle and cut the tip of the cartridge nozzle off. Replace the outer nozzle and cut its tip off so that a thread of gel approximately 5mm in diameter is able to be squeezed out. **VICTOR GEL** can now be dispensed as required in and around buildings, furniture and equipment.

## **APPLICATIONS RATES**

Bait application should consist of 0.25–0.5 g/spot and 10-20 spots/10 m<sup>2</sup>. A spot of 3 mm in diameter is equal to approximately 0.25 g of bait whereas a spot of 6 mm in diameter equals approximately 0.5 g of bait. The lower application rate can be used for low and moderate infestations or re-treatments, while the higher rate can be used for more severe infestations or a primary treatment.

Rapid control is provided by a number of small placements rather than less large spots. The gel will remain pliable and palatable to cockroaches as long as it is visible; usually for up to 20 weeks. If infestations are high, inspect the gel after 4 weeks of treatment and replenish if necessary. Daylight conditions will reduce the longevity of the bait.

**VICTOR GEL** must not be contaminated with other insecticides or placed on freshly sprayed surfaces, however, it can be integrated into other space or surface spray treatments.

## WAARSKUWINGS

- Hanteer versigtig.
- Skadelik indien ingesluk word.
- Skadelik vir visse en ander diere.
- Bewaar in 'n koel plek weg van voedsel en voer.
- Hou buite bereik van kinders, oningeligte persone en diere.

## Voorsorgmaatreëls

- Verhoed kontak deur mond of met vel.
- Dra rubberhandskoene tydens toediening.
- Was met water en seep na gebruik of na toevallige velkontak.
- Moenie eet, drink of rook tydens toediening of voordat hande en gesig gewas is nie.
- Dien toe op plekke wat totaal onbereikbaar is vir kinders en diere.
- Moenie damme, rioolstelsels, afvoerpype, riviere, damme of ander waterweë met die produk of gebruikte houers besoedel nie.
- Moet nie toedien op oppervlakke waar voedsel of voer geberg, voorberei of geëet word nie.
- Voorkom besoedeling van enige voedsel voorbereidings toerusting en eetgerei tydens toediening.
- Vernietig leë houers en moet nie vir geen ander doel gebruik nie.

## GEBRUIKSAANWYSINGS

### Gebruik slegs soos aangedui.

Geskik vir die gebruik in huise, restaurante, hotelle, hospitale, kantore, voedselverwerking- en vervaardigings aanlegte, handel- en nywerheidspersone, skole, skepe, stoorkamers, pakhuse, markplekke ens.

**VICTOR GEL** bevat 'n insekdoder met 'n vertraagde aksie, daarom is beheer nie onmiddelik sigbaar nie. 'n Sigbare effek op die kakkerlakpopulasie kan binne 'n week na behandeling verwag word. Ondersoek die area wat behandel moet word en identifiseer die kakkerlak infestasiëpunte.

**VICTOR GEL** word aangewend as 'n kolbehandeling in- en rondom kakkerlak skuilplekke. Die jel moet buite sig in donker plekke aangewend word.

**VICTOR GEL** word met 'n meganiese toediener aangewend soos volg:

Plaas die houers in die toediener raamwerk. Trek die sneller totdat die drukker kontak maak met die suier van die houers, sonder om druk toe te pas. Verwyder die vasgehegte spuitstuk en knip die voorste gedeelte van die houers se punt af. Plaas die vasgehegte spuitstuk terug en sny die punt af sodat 'n jelsrook, 5mm in deursnee uitgelaat kan word. **VICTOR GEL** kan nou toegedien word soos benodig word in en om geboue, toerusting, meubels, elektroniese, meganiese toerusting, ens.

## **TOEDIENINGSHOEVEELHEDE**

Gebruik die laer toedienings vir lae en gemiddelde besmettings of tydens opvolgbehandelings. Gebruik die hoër toedienings vir swaar besmettings of waar die eerste behandeling gedoen word. Aas toedienings moet min or meer gelyk wees aan 0.25–0.5 g/spikkel en 10–20 spikkels/10 m<sup>2</sup>. 'n Spikkel van ongeveer 3 mm in deursnee is gelykstaande aan 0.25 g aas. 'n Spikkel van ongeveer 6 mm in deursnee is gelykstaande aan 0.5 g aas.

Talryke kleiner uitplasings sal vinniger beheer gee as 'n paar groter spikkels. Solank die jel sigbaar is sal dit buigsaam en smaaklik bly vir kakkerlakke. Waar swaar besmettings voorkom, moet die jel vier weke na toediening ondersoek en vervang word indien nodig. Onder daglig-toestande mag die jel afgebreek word.

**VICTOR GEL** kan geïntegreer word met ander insekdoders, maar moet nie besoedel word met hierdie insekdoders of op vars bespuite oppervlakke geplaas word nie.