



# MAVRIK®



ADAMA



INSECTICIDES

Oil in water  
emulsion

MAPP  
10612

I



## Ingredients

- Tau-fluvalinate:  
240 g/l (22.0% w/w)



## Crops

A contact insecticide for the control of aphids in winter and spring wheat and winter and spring barley and control of pollen beetles, cabbage seed weevil and aphids in winter and spring oilseed rape and reduction in cabbage stem flea beetle in winter oilseed rape.



## Scan for Safety Data Sheet

Or use the weblink :  
<https://bit.ly/3QkgrYg>  
Alternatively, contact your supplier.



TRACEABIL-  
ITY PRINTING  
51/58 x 27 mm

1Le

**PROTECT FROM FROST SHAKE WELL BEFORE USE  
SAFETY PRECAUTIONS**

Open here 



**Warning**

**Very toxic to aquatic life with long lasting effects.**

Keep out of reach of children.  
Dispose of contents/container to a licensed hazardous-waste disposal contractor or collection site except for empty, clean containers which can be disposed of as non-hazardous waste.

**To avoid risks to human health and the environment, comply with the instructions for use.**

**Operator Protection**

*Engineering control of operator exposure must be used where reasonably practicable in addition to the following personal protective equipment:*

**WEAR SUITABLE PROTECTIVE CLOTHING (COVERALLS), SUITABLE PROTECTIVE GLOVES AND FACE PROTECTION (FACESHIELD)** when handling the concentrate.

**WEAR SUITABLE PROTECTIVE CLOTHING (COVERALLS) AND SUITABLE PROTECTIVE GLOVES** when handling contaminated surfaces.

**WEAR SUITABLE PROTECTIVE CLOTHING (COVERALLS)** when applying the product.

*However, engineering controls may replace personal protective equipment if a COSHH assessment shows they provide an equal or higher level of control.*

**AFTER CONTACT WITH SKIN, WASH IMMEDIATELY** with plenty of water.

**IN CASE OF CONTACT WITH EYES, RINSE IMMEDIATELY** with plenty of water and seek medical advice.

**DO NOT BREATHE SPRAY.**

**WASH HANDS AND EXPOSED SKIN** before meals and after work

**Environmental Protection**

Do not contaminate water with the product or its container. Do not clean application equipment near surface water. Avoid contamination via drains from farmyards and roads.

L1226421PWB\_01

**IMPORTANT INFORMATION**

FOR PROFESSIONAL USE ONLY AS AN AGRICULTURAL FUNGICIDE

**Crops:** Winter & spring wheat, winter & spring barley, winter & spring oilseed rape.  
**Maximum individual dose; Maximum total dose; Maximum number of treatments; Latest time of application and Other specific restrictions:** full details are given in the information box within the leaflet attached to the product.

**READ THE LABEL BEFORE USE. USING THIS PRODUCT IN A MANNER THAT IS INCONSISTENT WITH THE LABEL MAY BE AN OFFENCE. FOLLOW THE CODE OF PRACTICE FOR USING PLANT PROTECTION PRODUCTS.**

**This leaflet is part of the approved label.**

MAVRIK® is an oil in water emulsion containing 240 g/litre (22.0% w/w) tau-fluvalinate. It is a contact insecticide for the control of aphids in winter and spring wheat and winter and spring barley and control of pollen beetles, cabbage seed weevil and aphids in winter and spring oilseed rape and reduction in cabbage stem flea beetle in winter oilseed rape

---

**Environmental Protection continued.**

DO NOT ALLOW DIRECT SPRAY from horizontal boom sprayers to fall within 5m of the top of the bank of any static or flowing water body, or 1m of the top of a ditch which is dry at the time of application. DO NOT ALLOW DIRECT SPRAY from hand-held sprayers to fall within 1m of the top of the bank of a static or flowing waterbody. Aim spray away from water.

THIS PRODUCT IS NOT ELIGIBLE FOR BUFFER ZONE REDUCTION UNDER THE LERAP SCHEME.

HIGH RISK TO CERTAIN NON-TARGET INSECTS OR OTHER ARTHROPODS.

DO NOT SPRAY CEREALS within 6m of the field boundary.

**Storage and Disposal**

KEEP IN ORIGINAL CONTAINER, tightly closed, in a safe place.

RINSE CONTAINER THOROUGHLY, by using an integrated pressure rinsing device or manually rinsing three times. Add washings to sprayer at time of filling and dispose of safely.

*The (COSHH) Control of Substances Hazardous to Health Regulations may apply to the use of this product at work.*

**For advice on medical emergencies, fires or major spills telephone the National Chemical Emergency Centre on 01865 407333.**

---

**IMPORTANT INFORMATION**  
FOR PROFESSIONAL USE ONLY AS AN AGRICULTURAL INSECTICIDE

<b>Crops</b>	<b>Maximum individual dose (litres product/ha)</b>	<b>Maximum total dose (litres product/ha/crop)</b>	<b>Maximum number of treatments</b>	<b>Latest time of application</b>
Wheat (winter)	0.2	0.4	-	Before kernel medium milk
Barley (winter)	0.2	0.4	-	Before caryopsis watery ripe
Wheat (spring)	0.15	-	1 per crop	Before kernel medium milk
Barley (spring)	0.15	-	1 per crop	Before caryopsis watery ripe
Oilseed rape	0.2	0.4	-	Up to and including the end of flowering

**Other specific restrictions:**

This product must not be applied to a cereal crop if any other product containing a pyrethroid has been applied to that crop after the start of ear emergence.

A minimum interval of 14 days must be observed between applications when applied to listed cereals.

**READ THE LABEL BEFORE USE. USING THIS PRODUCT IN A MANNER THAT IS INCONSISTENT WITH THE LABEL MAY BE AN OFFENCE. FOLLOW THE CODE OF PRACTICE FOR USING PLANT PROTECTION PRODUCTS.**

## DIRECTIONS FOR USE

**IMPORTANT:** This information is approved as part of the Product Label. All instructions within this section must be read carefully in order to obtain safe and successful use of this product.

## RESTRICTIONS

### Safety to non-target arthropods

When used as directed at 0.15 litres per hectare, MAVRIK presents a low risk to ground (carabid) beetles, rove (*staphylinid*) beetles and adult hoverflies (*syrphids*), though some reduction in numbers may occur. Laboratory studies indicate that reductions in fecundity and egg viability may occur.

Avoid spraying oilseed rape within 6 metres of the field boundary to reduce effects on certain non-target insects or other arthropods.

### Resistance

Control may be reduced where strains of pests resistant to tau-fluvalinate develop.

## PEST CONTROL

### Winter and spring barley, winter and spring wheat

Summer aphids

Autumn /winter aphids

### Winter and spring oilseed rape

Pollen (blossom) beetle

Aphids

Cabbage stem flea beetle (winter oilseed rape)

Cabbage seed weevil

## CROP SPECIFIC INFORMATION

### Winter and spring barley, winter and spring wheat (summer aphid control)

**Maximum individual dose:** 0.15 litres per hectare

As a guide, spray once when aphids appear on two thirds of the ears and numbers are increasing. Utilise local predictive warnings if available.

### Winter barley and winter wheat (autumn/winter applications)

**Maximum individual dose:** 0.2 litres per hectare

Treat according to specialist advice, normally:

High Risk Crops: drilled in September/early October, in virus prone areas or following grass or weedy stubble.

Spray mid-October. A repeat application should be made during late autumn/early winter if aphid activity persists.

Medium Risk Crops; drilled before mid-October, in virus prone areas.

Spray late October/ early November\*.

Low Risk Crops; drilled after mid-October and crops in non virus prone areas.

Follow seasonal warnings based on aphid monitoring in local area\*.

N.B. \* In mild conditions, later sprays may be worthwhile.

**Maximum total dose (all cereal uses)**

Winter wheat & barley - 0.4 litres product per hectare per crop

Spring wheat & barley - 0.15 litres product per hectare per crop

**Latest time of application (cereals)**

Winter and spring wheat: Before kernel medium milk

Winter and spring barley: Before caryopsis watery ripe

**Winter and Spring Oilseed Rape**

Pollen (blossom) beetle

**Maximum individual dose:** 0.2 litres per hectare

Optimum control of this pest is achieved by application at the green to yellow bud stage. Apply according to local threshold advice.

A repeat application may be made if pollen beetle activity persists.

Aphids

**Maximum individual dose:** 0.2 litres per hectare

Not recommended for use to control aphids on spring rape. Spray according to local advice based on monitoring.

A repeat application may be made if aphid activity persists.

Cabbage stem flea beetle

Maximum individual dose: 0.2 litres per hectare

Egg hatch can be from October to March but mostly takes place in late autumn and from mid-late January. Apply according to local threshold advice.

MAVRİK will reduce damage caused by cabbage stem flea beetle in winter oilseed rape.

For optimal results a repeat application is advisable if cabbage stem flea beetle activity persists.

Cabbage seed weevil

Maximum individual dose: 0.2 litres per hectare

Apply during flowering period when the numbers of pests reach the threshold, or according to local advice based on monitoring. This is usually between 20% pod set and the end of flowering. For spring varieties, make the first application at green to yellow bud stage. Make a repeat application during flowering as required.

### **Maximum total dose**

Winter oilseed rape - 0.4 litres product per hectare per crop

Spring oilseed rape - 0.4 litres product per hectare per crop

### **Latest time of application**

Winter and spring oilseed rape: up to and including end of flowering

### **APPLICATION**

Good spray cover of the target is essential. Apply through a conventional hydraulic sprayer.

Do not leave the spray liquid in the sprayer for long periods (i.e. during meals or overnight).

### **SPRAY VOLUME**

A minimum of 200 litres of water per hectare should be used for all applications. Increase this volume of water in order to obtain good penetration into dense crops.

### **SPRAY QUALITY**

Apply as a MEDIUM quality spray as defined by BCPC. A minimum pressure of 2-3 bar should be used.

### **MIXING AND SPRAYING**

Shake the container of MAVRIK thoroughly before opening. Add half the required volume of clean water to the spray tank. Add the recommended quantity of MAVRIK. Agitate whilst filling the tank to the required water volume and continue agitation during spraying. Wash out all spray equipment with water immediately after use.

Crops should not be re-entered until spray residues are dry.

---

## **COMPANY ADVISORY SECTION**

*This section is not part of the Product Label under the Plant Protection Products Regulations and provides additional advice on product use at the discretion of ADAMA Agricultural Solutions UK Ltd.*

### **DISCLAIMER/CONDITIONS OF SUPPLY**

The specified properties of our products and the mode of application stated on this label have been established on the basis of research and experience. Products conform to specification at the time of delivery but, as we exercise no control over their subsequent storage, handling, mixing or use or the weather conditions before, during and after application, all of which may affect the performance of the products, no responsibility or liability will be accepted by us or our re-sellers for any failure in performance, damage or injury to person or property whatsoever arising from the storage, handling, application or use of the products. These conditions cannot be varied by our staff or agents whether or not they supervise or assist in or make recommendations concerning the use of such products. We recommend you contact your dealer to request advice on the suitability of this product for any new and/or unusual growing methods or for new varieties not listed on this label.

ADAMA Agricultural Solutions UK Ltd  
Third Floor East  
1410 Arlington Business Park  
Theale  
Reading  
RG7 4SA

Tel: 01635 860555  
Technical Helpline: 01635 876622  
[www.adama.com](http://www.adama.com)  
Email: [ukenquiries@adama.com](mailto:ukenquiries@adama.com)

MAVRİK® is a registered trademark of a company of the ADAMA Group.

© ADAMA Agricultural Solutions UK Ltd