

Key Benefits

- Provides excellent annual meadow-grass as well as broad-leaved weed control
- A useful tank mix partner for black-grass control
- Tank mix flexibility

Crop Information

Crop	Max individual dose (L/ha)	Maximum number of treatments (per crop)	Latest time of application
Winter wheat, Winter barley	2.5	1	Before leaf sheath erect stage (GS 30)
Winter rye, Winter triticale	2.5	1	Up to 4 true leaves (GS 14)
Spring wheat, Spring barley	2.5	1	Before leaf sheath erect stage (GS 30)

Product Information

Active ingredients:

40g/L diffufenican 400g/L pendimethalin

Formulation:

Suspension Concentrate

Pack size: 10 litres

Application Information

Water volume: 150-200L/ha
Spray quality: Medium

Aquatic buffer: 6 metres

Weed Spectrum

Troca o postrarii				
Weed	Pre – emergence 2.5L/ha	Post emergence 2.0L/ha (up to 1 true leaf stage)	Post emergence 2.5L/ha (up to 1 true leaf stage)	
Annual meadow-grass	S	-	-	
Black - bindweed	S	MR	MS	
Cleavers	S	MS	S	
Common chickweed	S	S	S	
Common poppy	MS	S	S	
Field pansy	S	S	S	
Field penny-cress	S	-	-	
Mayweeds	S	MS	S	
Red deadnettle	S	S	S	
Redshank	S	S	S	
Shepherd's purse	S	S	S	
Loose silky - bent	S	MS	S	
Smooth sowthistle	MS	-	-	
Speedwell	S	S	S	
Volunteer oilseed rape*	S	-	-	

^{*}Deep germinating volunteer oilseed rape may not be controlled.