

GUSTO[®] 3

PCS 04959

A ready-for-use, pelleted bait for the reduction of slug and snail damage in agricultural and horticultural crops.
Contains metaldehyde 3% w/w. Also contains calcium hydroxide.

FOR PROFESSIONAL USE ONLY


For advice on medical emergencies, fires or major spills telephone the UK National Chemical Emergency Centre on +44 (0)1865 407333

STORE IN A COOL, DRY PLACE

Authorised to & marketed by:
Adama Agricultural Solutions UK Ltd
Third Floor East, 1410 Arlington Business Park
Theale, Reading RG7 4SA
Telephone: +44 (0)1635 860555
Technical Helpline: +44 (0)1635 876622
www.adama.com
ukenquiries@adama.com



Meta is an assurance of quality
Metaldehyde active ingredient.
All Adama Crop Solutions UK Ltd slug pellets contain
Metaldehyde from Lonza.



ADAMA



GUSTO 3 is packaged in high quality
MINrinse bags

- Fully recyclable - Accepted by all of the major recycling and waste disposal companies.
- Not classified as hazardous waste - Empty packaging can be treated as a triple rinsed container.
- Easy to store - Packs can be stacked safely due to the unique antislip technology.



LDPE

Please recycle
this packaging.



Slug pellets are pesticides

- Keep pesticides out of water. → If you can't spray don't pellet!
- Do not contaminate drains or watercourses when cleaning the applicator.



SAFETY INFORMATION

Danger

Causes serious eye damage.
Suspected of damaging fertility.
Keep out of reach of children.
Obtain special instructions before use.
Do not handle until all safety precautions have been read and understood.
Wear protective gloves/protective clothing/eye protection/face protection.
IF IN EYES: rinse cautiously with water for several minutes. Remove contact lenses if present and easy to do. Continue rinsing.
Immediately call a POISON CENTRE or doctor/physician.

Store locked up.
If medical advice is needed, have product container or label at hand.
Dispose of contents/container to a licensed hazardous waste disposal contractor or collection site except for empty, clean containers which can be disposed of as non-hazardous waste.
To avoid risks to human health and the environment, comply with the instructions for use.

PCS 04959

To access the Safety Data Sheet for this product scan the QR code or use the weblink:
<https://bit.ly/357m831>
Alternatively, contact your supplier.

Scan for
Safety Data
Sheet



IMPORTANT INFORMATION

FOR PROFESSIONAL USE ONLY AS AN AGRICULTURAL/HORTICULTURAL MOLLUSCICIDE

Crop	Maximum individual dose (kg product/ha)	Maximum total dose (kg product/ha)	Latest time of application
Potato	6	12	GS 97 (3 days before harvest)
Wheat, barley, oats, rye & triticale	6	12	GS 29 (60 days before harvest)
Strawberry (outdoor)	6	12	GS 41
Sugar and fodder beet	6	12	GS 19 (90 days before harvest)
Cauliflower, Brussels sprout, broccoli	6	12	GS 49 (7 days before harvest)
Beans, peas	6	12	GS 50 (21 days before harvest)
Oilseed rape	6	12	GS 19 (90 days before harvest)
Ornamentals	6	12	-

15 kg

WARNING: Keep secure at all times.
This product may be attractive to animals.

Batch No.: see packaging

R3.2 JAN 21



SAFETY PRECAUTIONS

Operator Protection

WHEN USING, DO NOT EAT, DRINK OR SMOKE.

WASH HANDS before eating and drinking after work.

Environmental Protection

Do not contaminate water with the product or its container.

Do not clean application equipment near surface water.

Avoid contamination via drains from farmyards and roads.

DANGEROUS TO GAME, WILD BIRDS AND ANIMALS. Bury or remove all spillages.

KEEP POULTRY out of treated areas for at least seven days after treatment.

Storage and Disposal

KEEP AWAY FROM FOOD, DRINK AND ANIMAL FEEDINGSTUFFS.

KEEP IN ORIGINAL CONTAINER, tightly closed, in a safe place.

EMPTY CONTAINER COMPLETELY and dispose of safely.

DIRECTIONS FOR USE

IMPORTANT: This information is approved as part of the Product Label. All instructions within this section must be read carefully in order to obtain safe and successful use of this product.

RATE OF APPLICATION

All crops

GUSTO® 3 is applied at 6 kg product/ha

The maximum total dose must not exceed 12 kg of product/ha/year.

In high risk situations, where additional treatments against slugs are necessary beyond the permissible 360 g metaldehyde/ha/year total dose, use an alternative slug control product not containing metaldehyde.

APPLICATION

Broadcast application

Apply using a fertiliser distributor or a purpose made granule applicator. Small areas may be treated by hand or hand-held equipment. Calibrate all equipment before use.

Crop	Timing/method
Cereals	Pre-drilling Apply to the prepared seed bed and leave 4-7 days before drilling. Post-drilling Broadcast as soon as damage is seen or indicated by test baiting.
Oilseed rape	Broadcast shortly after drilling.
Sugar beet Fodder beet	Broadcast evenly over the soil surface when slugs and snails are active preferably before crop damage occurs.
Other agricultural and horticultural crops	Broadcast evenly over the soil surface when slugs and snails are active preferably before crop damage occurs. See Note on 'late application to broad-leaved plants' under the heading 'Important' below.

Slug Trapping

To establish the need for pellet application on winter wheat or winter oilseed rape, monitor for slug activity. Where bait traps are used, use a foodstuff attractive to slugs e.g. chicken layers' mash which has proven to be particularly effective. DO NOT use slug pellets in traps since they are a potential hazard to wildlife and pets. Details of the correct baiting procedure are provided in the 'COMPANY ADVISORY INFORMATION' section of this label.

For further information on slug control and damage risk assessment, please refer to the UK AHDB Information Sheet 04 Winter 2018 (Integrated Slug Control) which can be found at:

www.ahdb.org.uk/knowledge-library/integrated-slug-control

or use the weblink: <http://bit.ly/sluginfo>

Important

- Complete control of slugs/snails is difficult to achieve with any type of molluscicide, so some crop loss may occur especially in wet seasons. Best control is achieved by a programme of good husbandry (see COMPANY ADVISORY INFORMATION) and properly timed applications of a pellet bait such as GUSTO 3.
- The first application of GUSTO 3 should be made as a preventative measure as specified in the APPLICATION table. Follow-up applications may be required, especially in wet seasons. Assessment by test baiting will indicate when slugs are active at the soil surface and if conditions are suitable for further GUSTO 3 treatment.
- Late application of pellets to broad-leaved plants may result in lodging of the pellets in foliage. Care should be taken to avoid this when making applications to edible crops, e.g. cauliflower. In cauliflower apply prior to hearting or curd formation.
- AVOID application if heavy rain is expected, as this may reduce effectiveness of the pellets.
- Best results are obtained when slugs and snails are actively feeding, generally following a period of light rain in mild conditions. Where a heavy attack is likely to occur, re-apply 2-3 weeks later or whenever test baiting shows it to be necessary.

DISCLAIMER/CONDITIONS OF SUPPLY

The specified properties of our products and the mode of application stated on this label have been established on the basis of research and experience. Products conform to specification at the time of delivery but, as we exercise no control over their subsequent storage, handling, mixing or use or the weather conditions before, during and after application, all of which may affect the performance of the products, no responsibility or liability will be accepted by us or our re-sellers for any failure in performance, damage or injury to person or property whatsoever arising from the storage, handling, application or use of the products. These conditions cannot be varied by our staff or agents whether or not they supervise or assist in or make recommendations concerning the use of such products. We recommend you contact your dealer to request advice on the suitability of this product for any new and/or unusual growing methods or for new varieties not listed on this label.

COMPANY ADVISORY INFORMATION

(Note: This section is not part of the approved Directions for Use)

GUSTO 3 SLUG AND SNAIL CONTROL PROGRAMME

Slugs and snails may be controlled by a programme of good husbandry and accurately timed applications of GUSTO 3 – see DIRECTIONS FOR USE.

STEP 1: GOOD HUSBANDRY

Observe and where possible, practise the following husbandry techniques to reduce the overall population of slugs and snails. Under very wet conditions or where the slug population is very high, a sufficient number of slugs may survive applications of GUSTO 3 to cause crop damage. Good husbandry is essential under such conditions to minimise crop damage.

Slug Pellets are pesticides, keep pesticides out of water.

- o Calibrate your applicator before use.
- o Fill and clean application equipment in the field.
- o Do not fill or clean down in the yard.
- Avoid applying slug pellets when heavy rain is forecast.
- Check that the product packaging is intact and clear up spillages immediately.
- If possible, avoid sowing slug susceptible crops after leafy crops, such as oilseed rape, which can carry high populations especially in wet seasons.
- Aim to minimise crop debris to minimise slug habitat in the field.
- Plough or cultivate under dry soil and weather conditions to reduce the slug population substantially.
- Prepare a fine seedbed with a firm tilth. Slugs do not burrow through the ground but follow cracks or openings in the soil.
- Mineral, light and drier soils generally carry lower slug populations than heavier soils. Irrigation favours the slugs.
- Slugs and snails when surface feeding, do so mainly at night, so inspect fields for them at dawn or dusk.

STEP 2: APPLICATIONS OF GUSTO 3

GUSTO 3 should be applied as part of an IPM programme for slug and snail control. The need for molluscicide application should be determined by assessing the risk of slug damage e.g. using a test baiting procedure. If molluscicide application is required, GUSTO 3 should be applied as specified in the DIRECTIONS FOR USE. Follow-up applications may be required, especially in wet seasons. The timing of further applications must be established by test baiting. If the maximum application rate of metaldehyde-containing slug pellets is reached, control via an alternative measure should be used. Some species of slugs may live and feed underground for long periods without coming to the surface. Only slugs which surface at, or soon after application, have the opportunity to feed on the pellets.

Test Baiting Procedure

- Put slug traps out before cultivation, when the soil surface is visibly moist and the weather mild (5-25°C).
- Place upturned flower pot bases about 25 cm diameter with a small heap (20 ml or 2 heaped teaspoonfuls of chicken layers' mash (NOT slug pellets) beneath each trap.
- Position these traps on the areas of the field where slug damage has been historically bad. Ideally 9 slug traps in a W pattern or 13 if the field is larger than 20 ha.
- Traps should be left overnight and examined the following morning.
 - FOR WINTER WHEAT, a catch of 4 or more slugs per trap indicates a possible risk, where soil and weather conditions favour slug activity.
 - FOR WINTER OILSEED RAPE a catch of 4 or more slugs per trap in standing cereals or 1 or more in cereal stubble, if other conditions were met, would also indicate a possible risk of damage.

GUSTO® is a registered trademark of the Adama Group.
© Adama Agricultural Solutions UK Ltd

GUSTO[®] 3

PCS 04959

A ready-for-use, pelleted bait for the reduction of slug and snail damage in agricultural and horticultural crops.
Contains metaldehyde 3% w/w. Also contains calcium hydroxide.

FOR PROFESSIONAL USE ONLY


For advice on medical emergencies, fires or major spills telephone the UK National Chemical Emergency Centre on +44 (0)1865 407333

STORE IN A COOL, DRY PLACE

Authorised to & marketed by:
Adama Agricultural Solutions UK Ltd
Third Floor East, 1410 Arlington Business Park
Theale, Reading RG7 4SA
Telephone: +44 (0)1635 860555
Technical Helpline: +44 (0)1635 876622
www.adama.com
ukenquiries@adama.com



Meta is an assurance of quality
Metaldehyde active ingredient.
All Adama Crop Solutions UK Ltd slug pellets contain
Metaldehyde from Lonza.



ADAMA



GUSTO 3 is packaged in high quality
MINrinse bags

- Fully recyclable - Accepted by all of the major recycling and waste disposal companies.
- Not classified as hazardous waste - Empty packaging can be treated as a triple rinsed container.
- Easy to store - Packs can be stacked safely due to the unique antislip technology.



LDPE

Please recycle
this packaging.



Slug pellets are pesticides

- Keep pesticides out of water. → If you can't spray don't pellet!
- Do not contaminate drains or watercourses when cleaning the applicator.



SAFETY INFORMATION

Danger

Causes serious eye damage.
Suspected of damaging fertility.
Keep out of reach of children.
Obtain special instructions before use.
Do not handle until all safety precautions have been read and understood.
Wear protective gloves/protective clothing/eye protection/face protection.
IF IN EYES: rinse cautiously with water for several minutes. Remove contact lenses if present and easy to do. Continue rinsing.
Immediately call a POISON CENTRE or doctor/physician.

Store locked up.
If medical advice is needed, have product container or label at hand.
Dispose of contents/container to a licensed hazardous waste disposal contractor or collection site except for empty, clean containers which can be disposed of as non-hazardous waste.
To avoid risks to human health and the environment, comply with the instructions for use.

PCS 04959

To access the Safety Data Sheet for this product scan the QR code or use the weblink:
<https://bit.ly/357m831>
Alternatively, contact your supplier.

Scan for
Safety Data
Sheet



IMPORTANT INFORMATION

FOR PROFESSIONAL USE ONLY AS AN AGRICULTURAL/HORTICULTURAL MOLLUSCICIDE

Crop	Maximum individual dose (kg product/ha)	Maximum total dose (kg product/ha)	Latest time of application
Potato	6	12	GS 97 (3 days before harvest)
Wheat, barley, oats, rye & triticale	6	12	GS 29 (60 days before harvest)
Strawberry (outdoor)	6	12	GS 41
Sugar and fodder beet	6	12	GS 19 (90 days before harvest)
Cauliflower, Brussels sprout, broccoli	6	12	GS 49 (7 days before harvest)
Beans, peas	6	12	GS 50 (21 days before harvest)
Oilseed rape	6	12	GS 19 (90 days before harvest)
Ornamentals	6	12	-

15 kg

WARNING: Keep secure at all times.
This product may be attractive to animals.

Batch No.: see packaging

R3.2 JAN 21