


FirstAct™ Herbicide for the Double Team Sorghum Cropping Solution

FirstAct herbicide delivers Double Team Sorghum growers reliable broad spectrum grass control along with flexible rotation and high yields for more profitable returns per acre.

Benefits

- High yield potential and protection
- Improved weed control and strong post-emergence grass control
- Crop safety for Double Team Sorghum
- Convenient application





Use Directions

- Do not apply FirstAct before the 8-leaf growth stage (approx. 11-12" sorghum) or significant crop injury may occur
- If required, an over-the-top broadleaf herbicide spray should be made at least 7 days prior to FirstAct application
- Apply 5-12 fl oz/acre (most FirstAct rates will be 8 – 10 oz/acre)
- Difficult-to-control perennial weeds will require a use rate of 12 oz/acre
- Recommended minimum carrier volume of 15 gal/acre.
- Add crop oil concentrate (COC) at 1% v/v
- Two applications are allowed with maximum total of 21 oz/acre/year
- FirstAct has a 45-day pre-harvest interval (PHI)

Ground Application Parameters

- Use spray nozzles that will deliver medium or larger spray droplets
- Do not use flood, rain drop, whirl chamber, or other nozzle types that produce coarse, large spray droplets
- Use a minimum of 10 gallons of water per acre or a minimum of 15 gallons of water per acre in arid areas
- Increase spray volume and pressure as weed or crop density and size increase

Aerial Application Parameters

- Use spray nozzles that will deliver coarse or larger spray droplets
- Use nozzle types and arrangements that provide optimum spray distribution and maximum coverage
- Use a minimum of 3 gallons of water per acre or a minimum of 5 gallons of water per acre in arid areas

Mixing Instructions

1. Fill the tank $\frac{1}{4}$ to $\frac{1}{3}$ full of water.
2. While agitating, add the required amount of FirstAct.
3. Continue agitation until the FirstAct is fully dispersed, at least 5 minutes.
4. Once the FirstAct is fully dispersed, maintain agitation and continue filling tank with water.
5. As the tank is filling, add the required volume of spray additives, always add these to the spray tank last.
6. Apply FirstAct spray mixture within a reasonable period of time of mixing to avoid product degradation (24 to 48 hours). If the spray mixture stands for any period of time, thoroughly re-agitate before using.

Crop Rotations

Do not rotate to crops other than the following within 120 days after application:

- | | | |
|-----------------------------------|----------------------------------|----------------------------------|
| · Cotton | · Rapeseed Crop Subgroup 20A | · Snap Beans |
| · Dry Beans (including Chickpea) | [includes Borage, Canola, | · Soybeans |
| · Flax | Crambe, Gold of Pleasure | · Sunflowers |
| · Lentils | (Camelina), Cuphea, Echium, | · Sugarbeets |
| · Mint (Spearmint and Peppermint) | Hare's Ear Mustard, Lesquerella, | · Quizalofop-tolerant field corn |
| · Peas (Dry and Succulent Peas) | Lunaria, Meadowfoam, Milkweed, | |
| | Mustard Seed, Oil Radish, Poppy | |
| | Seed, Sesame, and Sweet Rocket] | |

Target Weeds

Volunteer Corn, Sandbur,
Texas Millet/Panicum,
Crabgrass, Foxtail and
Barnyardgrass*

**See label for additional grasses*

Mode of Action

ACCase inhibitor
(WSSA Group 1)

Signal Word

Caution

Restricted Use

No

