

Active Ingredients:

Tebuconazole 200 g/L and Azoxystrobin 120 g/L = SC

Application Rates and Acres Treated:

Rate: 190 - 250 ml/ac

Acres Treated: 40 - 53 ac/jug

Packaging:

Case: 2 × 10.08 L jugs

Water Volume:

Ground: 40 L/ac (10 US gal/ac)

Aerial: 20 L/ac (5 US gal/ac)

Rainfastness:

Avoid application when heavy rain is forecast.

KEY BENEFITS:

- Multiple modes of action for resistance management
- Curative and preventative disease control
- Broad spectrum leaf disease control with superior rust control compared to competitors
- · Penetrates quickly into the leaf tissue
- Keeps the plant greener longer

KEY DISEASES CONTROLLED:

- Crown rust
- Leaf rust
- Stem rust
- Stripe rust


- Septoria leaf blotch
- Tan spot
- Net blotch
- Spot blotch


HOW IT WORKS:


CUSTODIA® is a broad spectrum fungicide with both preventative and curative properties. With its unique systemic and translaminar activity, CUSTODIA® has shown to both move through the leaves and down the plant for more protection. CUSTODIA® penetrates quickly into the leaf tissue protecting the flag leaf from all major leaf diseases.

CUSTODIA®

Field Test Results








REGISTERED CROPS AND STAGING:

Crop	Diseases	Application Timing	Rate
Wheat (spring, winter, durum)	Leaf rust, Stem rust, Stripe rust, Septoria leaf blotch, Tan spot	Apply CUSTODIA® to leaf foliage at the first sign or very early stage of disease, up to the beginning of heading.	190 – 250 ml/ac
		Use of the higher rate should be considered when weather conditions are conducive to heavy disease development.	
Barley	Net blotch, Spot blotch, Leaf rust, Stem rust, Stripe rust, Septoria leaf blotch, Tan spot	Apply CUSTODIA® to leaf foliage at the first sign or very early stage of disease, up to the beginning of heading.	190 – 250 ml/ac
		Use of the higher rate should be considered when weather conditions are conducive to heavy disease development.	
Oats	Crown rust, Stem rust, Septoria leaf blotch	Apply CUSTODIA® foliar fungicide at the very early stages of disease development.	190 ml/ac
Soybeans	Asian soybean rust, Frogeye leaf spot	Apply CUSTODIA® foliar fungicide at the very early stages of disease development. Use of the higher rate should be considered when weather conditions are conducive to heavy disease development or when heavy disease pressure is present.	190 - 250 ml/ac

CUSTODIA®

Orchestrated Leaf Disease Control





CUSTODIA®

MIXING INSTRUCTIONS:

- 1. Fill the clean spray tank ¾ full with clean water.
- Add the required amount of CUSTODIA® Foliar Fungicide into the sprayer and agitate thoroughly.
- Continue agitation and add the required amount of the tank-mix partner.
- 4. Complete filling the tank to the desired level with water.

CROP ROTATIONS:

No restrictions.

PRE-HARVEST INTERVALS:

Mature grains: 45 daysForage, hay: 36 days

GRAZING RESTRICTIONS:

Do not allow livestock to graze or feed green forage to livestock prior to 6 days after treatment. Straw cut after harvest may be fed or used for bedding.

STORAGE:

Do not freeze.

Quick Tips

CUSTODIA® should be applied at flag leaf for optimal leaf disease control.

Coverage is key; do not use less than recommended water volumes.

For fusarium control, we would recommend an application of ORIUS® 430 SC fungicide at full head emergence after CUSTODIA® at flag leaf.

For resistance management, CUSTODIA® contains Group 3 and 11 fungicides.

When possible, rotate the use of CUSTODIA® or other Group 3 and 11 fungicides with different groups that control the same pathogens.



Always read and follow label directions. Toll-free: 1.855.264.6262 Website: ADAMA.COM

BUMPER®, CUSTODIA® and ORIUS® are registered trademarks of ADAMA Agricultural Solutions Canada Ltd. All other products are trademarks of their respective companies. © 2023 ADAMA Agricultural Solutions Canada Ltd.