

Active Ingredients:

Clodinafop-propargyl

Herbicide Group:

Group 1

Formulation:

Clodinafop-propargyl 80 g/L with built-in surfactant = EC

Application Rates & Acres Treated:

Rate: 283-356 ml/ac Acres Treated: 15-20 ac/jug,

255-320 ac/drum

Packaging:

Case: 2 × 5.66 L jugs
Pallet: 5 × 90.6 L drums

REGISTERED CROPS AND STAGING:

Wheat (spring, durum)	Prior to emergence of 4th tiller

WEEDS CONTROLLED:

WEED CONTROLLED	WEED STAGE	ADDITIONAL INFORMATION
Barnyard grass	1–5 leaf stage on main stem	For optimum control apply before tillering, and while barnyard grass is actively growing
Foxtail (green, yellow)	1–5 leaf stage on main stem	For optimum control apply prior to emergence of the 3 rd tiller and while foxtail is actively growing
Persian darnel	1–5 leaf stage on main stem	For optimum control apply before tillering, and while Persian darnel is actively growing
Voluntary canary seed	1–6 leaf stage on main stem	Prior to emergence of 4 th tiller
Volunteer oats (tame)	3–6 leaf stage on main stem	Prior to emergence of 4 th tiller
Wild oats	1–6 leaf stage on main stem	Prior to emergence of 4 th tiller

HOW IT WORKS:

LADDER ALL IN® is absorbed by the leaves and is rapidly translocated to the growing points of leaves and stems. Thorough coverage of the plants is essential for consistent control. Actively growing susceptible grasses stop growing within 48 hours of treatment. Depending on species, growing conditions and crop competition, leaves and growing points turn yellow within 1–3 weeks after application. Further colour changes and loss of vigour will be observed, followed by a browning and complete control 3–5 weeks after application.



LADDER ALL IN®

WATER VOLUME:

• Ground: 5-10 US gal/ac

Aerial: 3 US gal/ac

RAINFASTNESS:

30 minutes

REGISTERED AND SUPPORTED TANK MIXES:

Herbicides:

2,4-D amine

Ally®

BADGE®

BROMOTRIL®

Curtail[®] M

Dicamba

Dichlorprop-DX

Estaprop® XT

ESTEEM®

FORCEFIGHTER® M

Infinity®

Lontrel™ XC

MCPA amine

MCPA ester

MCPA sodium 300

· Mecoprop-P

OUTSHINE®1

Pixxaro[®]

• Pulsar®

• Refine® SG

Retain®SG

RUSH® 24

Target®

• THRASHER®

Trophy[®]

Insecticides:

• Decis®

SILENCER® 120 EC

Fungicides:

BUMPER® 432 EC

¹Normal weed pressure use with high rate of LADDER ALL IN® low rate only with low populations and early application.

When tank mixing with a broadleaf herbicide, insecticide, or a fungicide, always refer to the label of the broadleaf partner, insecticide partner or fungicide partner prior to use.

MIXING INSTRUCTIONS:

- 1. Clean spray tank and ½ fill with clean water. Start agitation or bypass system.
- 2. If a broadleaf herbicide, insecticide or fungicide is to be used, add the product FIRST prior to adding LADDER ALL IN® and agitate for 2-3 minutes.
- 3. Add correct amount of LADDER ALL IN®. Agitate for 3 5 minutes before adding remainder of water and then maintain constant agitation.
- 4. After any break in spraying operations, agitate thoroughly before spraying again.
- 5. Use the spray suspension as soon as it is prepared.
- 6. If an oil film starts to build up in the tank, drain tank and then clean with a detergent.

PRE-HARVEST INTERVAL:

60 days

GRAZING RESTRICTIONS:

Observe a minimum of 3 days before grazing livestock on treated crops.

STORAGE:

May be stored at any temperature.

Quick Tips

- - LADDER ALL IN® contains an



Always read and follow label directions. Toll-free: 1.855.264.6262 Website: ADAMA.COM/CANADA

BADGE®, BROMOTRIL®, BUMPER®, ESTEEM®, FORCEFIGHTER®, LADDER ALL IN®, OUTSHINE®, RUSH®, SILENCER®, THRASHER® and TOPNOTCH® are registered trademarks, and BRAZEN(TM) are trademarks of ADAMA Agricultural Solutions Canada Ltd. © 2021 ADAMA Agricultural Solutions Canada Ltd.