



ADAMA

# BADGE®

Multi-mode of action tank-mix partner  
for cereals, flax and corn.



## GROUP 4 & 6

### Active Ingredients:

Bromoxynil 225 g/L and MCPA  
(as 2-EH Ester) 225 g/L = EC

### Application Rates and Acres Treated:

Rate: 500 ml/ac

Acres Treated: 20 ac/jug;  
240 ac/drum; 900 ac/tote

### Packaging:

Case: 2 × 10 L jugs  
Bulk: 120 L drums  
Tote: 450 L

### Water Volume:

Ground: 20–40 L/ac (5–10 US gal/ac)  
in cereals and flax; 80–120 L/ac  
(20–30 US gal/ac) in corn; 60 L/ac  
(15 US gal/ac) in forages

Aerial (wheat, barley and oats only):  
8–20 L/ac (2–5 US gal/ac)

### Rainfastness:

1 hour

### REGISTERED CROPS:

- Barley
- Canary seed
- Corn
- Fall rye
- Flax
- Oats
- Seedling grasses
- Timothy (established for seed production)
- Wheat (spring, winter, durum)

### WEEDS CONTROLLED:

- American nightshade
- Ball mustard
- Bluebur
- Canada thistle<sup>1</sup>
- Cocklebur
- Common buckwheat
- Common groundsel
- Common ragweed
- Cow cockle<sup>2</sup>
- Flixweed
- Green smartweed
- Kochia<sup>3</sup>
- Lady's thumb
- Lamb's quarters
- Night-flowering catchfly
- Pale smartweed
- Perennial sow thistle<sup>1</sup>
- Redroot pigweed
- Russian thistle<sup>3</sup>
- Scentless chamomile<sup>4</sup>
- Shepherd's purse
- Stinkweed
- Tartary buckwheat
- Velvetleaf<sup>5</sup>
- Volunteer canola (all types)
- Volunteer sunflower
- Wild buckwheat
- Wild mustard
- Wild tomato
- Wormseed mustard

<sup>1</sup>Top growth control.

<sup>2</sup>Up to 4-leaf stage.

<sup>3</sup>Spray before plants are 2 inches high.

<sup>4</sup>Spring annual only.

<sup>5</sup>Spray before plants are 3 inches high.



# BADGE®

## HOW IT WORKS:

A combination of systemic and contact activity with weeds yellowing within 2–4 days and exhibiting abnormal growth (twisting and cupping of leaves) in 2–10 days.

## CROP STAGING:

Crop	Stage
Barley, Oats, Wheat (spring, durum)	2 leaf to early flag
Canary seed	3–5 leaf
Corn	4–6 leaf
Fall rye	When growth commences in spring to early flag leaf
Flax (including low linolenic varieties)	2 inches to early bud stage. Best tolerance occurs when flax is 2–4 inches tall.
Seedling grasses	2–4 leaf
Timothy (established for seed production)	Prior to shot blade in the seed production year
Winter wheat	2–4 leaf stage in the fall or after growth resumes up to early flag leaf

## REGISTERED AND SUPPORTED TANK MIXES:

- Corn: Atrazine
- Flax: ARROW® 240 EC, ARROW ALL IN® or Poast®
- Oats: MCPA ester
- Spring wheat and barley: Ally®, MCPA Ester, Refine® SG, BISON® 400 L
- Spring wheat only: LADDER ALL IN®, Everest® 3.0 or Traxos®
- Winter wheat: Refine® SG or Everest® 3.0

## MIXING INSTRUCTIONS:

1. Fill clean spray tank ½ full with water.
2. Add the required amount of BADGE® and agitate thoroughly.
3. Fill the tank and agitate again before use.
4. If tank mixing with a grassy weed herbicide, read both labels and follow the more stringent directions for tank mixing.

## PRE-HARVEST INTERVAL:

Flax: 60 days

## STORAGE:

Do not freeze.

## GRAZING RESTRICTIONS:

Do not graze treated grain or established timothy crops or cut for feed within 30 days of application.

## CROP ROTATIONS:

No re-cropping restrictions the year after treatment.

### QUICK TIPS:

Avoid spraying if temperatures are above 25° C.



ADAMA

Always read and follow label directions.  
Toll-free: 1.855.264.6262  
Website: ADAMA.COM

ARROW®, ARROW ALL IN®, BADGE®, BISON® and LADDER ALL IN® are registered trademarks of ADAMA Agricultural Solutions Canada Ltd. All other products are trademarks of their respective companies. © 2023 ADAMA Agricultural Solutions Canada Ltd.