

Active Ingredients:

Fluroxypyr 180 g/L, Bromoxynil 225 g/L, MCPA 2 EH Ester 225 g/L = EC

Application Rates and Acres Treated:

Rate:

500 ml/ac Bromoxynil/MCPA Ester; 240 ml/ac Fluroxypyr

Acres Treated:

40 ac/case; 480 ac/pallet

Packaging:

Case:

Bromoxynil/MCPA Ester: 2 × 10 L jugs; Fluroxypyr: 1 × 9.6 L jug

Pallet:

Bromoxynil/MCPA Ester: 2 × 120 L drums; Fluroxypyr: 1 × 115.2 L drum

Water Volume:

Ground: 20-40 L/ac (5-10 US gal/ac)Aerial: Do not apply.

Rainfastness:

1 hour

GROUP 4 & 6

REGISTERED CROPS: Barley

WEEDS CONTROLLED:

- · American nightshade
- Bluebur
- Burdock
- Canada thistle¹
- Chickweed*
- Cleavers (including Group 2 resistant biotypes)
- Cocklebur
- Common groundsel
- Cow cockle
- Flixweed
- Hempnettle*
- Kochia (including Group 2 and glyphosate-resistant biotypes)
- Lady's thumb

*Suppression only.

¹Top growth control.

- Wheat (spring, durum)
- · Lamb's quarters
- Night-flowering catchfly
- · Perennial sow thistle
- · Redroot pigweed
- Russian thistle
- Shepherd's purse
- Smartweed
- Stinkweed Vetch
- Volunteer canola
- Volunteer sunflower
- Wild buckwheat
- · Wild mustard (including Group 2 resistant biotypes)
- Wild radish
- Wormseed mustard

HOW IT WORKS:

Quickly causes plants to stop growing. Convenient co-pack controls a wide range of weeds, including glyphosateresistant and Group 2 resistant kochia, Group 2 resistant cleavers and Group 2 resistant wild mustard.





FORCEFIGHTER® M

CROP STAGING:

Crop	Stage
Wheat	2 leaf to early flag leaf
Barley	2 leaf to early flag leaf

REGISTERED AND SUPPORTED TANK MIXES:

Spring wheat:

- BISON® 400 L
- BRAZEN™ II
- Everest® 3.0
- LADDER ALL IN®
- Simplicity® GoDRI
- Refine® SG
- Traxos®

Durum wheat:

- Everest® 3.0
- LADDER ALL IN®
- · Simplicity® GoDRI
- Traxos®

Barley:

- BISON® 400 L
- BRAZEN™ II

MIXING INSTRUCTIONS:

- 1. Fill spray tank ½ full with water.
- 2. Add the required amount of FORCEFIGHTER® M and agitate thoroughly.
- 3. Add any tank-mix partners.
- 4. Fill the tank and agitate again before use.

CROP ROTATIONS:

Can be seeded the following year to barley, canola, flax, forage grasses, lentils, mustard, field peas, rye, and wheat.

GRAZING RESTRICTIONS:

30 days

STORAGE:

Do not freeze.

PRE-HARVEST INTERVAL:

60 days

QUICK TIPS:

FORCEFIGHTER® M's activity is influenced by weather conditions. The optimal temperature is 12 – 24° C. Avoid application 3 days before or after frost. Do not apply before the 2-leaf stage as crop injury may occur. Use 38 L/ac application volume when there is a heavy canopy or when most weeds are at an advanced stage of growth. Do not apply by air.

