

Active Ingredient:

Pinoxaden 50 g/L = EC

Application Rates & Acres Treated:

Rate: 500 ml/ac

Acres Treated: 20 ac/jug

Packaging:

Case: 2 x 10 L jugs

Water Volume:

Ground: 20 - 40 L/ac

(5-10 US gal/ac)

Aerial: 12 L/ac

(3 US gal/ac)

Rainfastness:

1 hour

GROUP 1

REGISTERED CROPS:

Barley

WEEDS CONTROLLED:

- Barnyard Grass
- Green & Yellow Foxtail
- Proso Millet

- Wheat (spring, winter)
- Volunteer Canary Seed
- Volunteer Oats
- Wild Oats

HOW IT WORKS:

BRAZEN ALL IN™ Herbicide is absorbed by the leaves and is rapidly translocated to the growing points of leaves and stems. Thorough coverage of the plants is essential for consistent control. Actively growing susceptible grasses stop growing within 48 hours of treatment. Depending on species, growing conditions and crop competition, leaves and growing points turn yellow within one to three weeks after application. Further colour changes and loss of vigour will be observed, followed by a browning and control three to five weeks after application.

CROP/WEED STAGING:

Crops/Weeds	Growth Stage
Wheat (spring, winter), Barley	1 leaf to flag leaf stage
Wild Oats, Green and Yellow Foxtail, Barnyard Grass, Volunteer Oats, Volunteer Canary Seed, Proso Millet	1–6 leaf, prior to 4 th tiller

^{*}Do not apply past flag leaf stage

CROP ROTATIONS: GRAZING RESTRICTIONS:

No restrictions. 7 days

PRE-HARVEST INTERVALS: STORAGE:

Grain, Straw: 60 days If frozen, allow to thaw and Hay: 30 days agitate thoroughly prior to use.



BRAZEN ALL IN™

REGISTERED AND SUPPORTED TANK MIXES¹

Herbicides:

- BADGE®
- Barricade® II
- BROMOTRIL®
- Cirpreme[™] XC
- Curtail® M2
- ESTEEM®
- ESTEEM ALL IN™

- Exhilarate™
- FORCEFIGHTER® M
- FORCEFIGHTER ALL IN™
- Infinity^{®2}
- Infinity® FX
- MCPA Ester 600²
- MCPA Amine (assume 500 series)
- OUTSHINE®

- OUTSHINE ALL IN™
- Pixxaro™
- Pulsar®
- Refine® SG
- TOPLINE®
- Travallas®
- Trophy®2

Fungicides:

• BUMPER® 432 EC

TOPNOTCH[™]

This product may be tank mixed with a fertilizer, a supplement, or with registered pest control products, whose labels also allow tank mixing, provided the entirety of both labels, including Directions For Use, Precautions, Restrictions, Environmental Precautions, and Spray Buffer Zones are followed for each product. In cases where these requirements differ between the tank mix partner labels, the most restrictive label must be followed. Do not tank mix products containing the same active ingredient unless specifically listed on this label.

In some cases, tank mixing pest control products can result in reduced pesticide efficacy or increased host crop injury. The user should contact ADAMA Agricultural Solutions Canada Ltd. at 1-855-264-6262 for information before applying any tank mix that is not specifically recommended on this label.

MIXING INSTRUCTIONS:

- 1. Ensure that the sprayer interior is clean, then fill the spray tank with ½ the required amount of water and engage gentle agitation. Good agitation is indicated by a rippling or rolling action on the surface
- 2. Add any WG or DF formulation mix partners and agitate to ensure complete mixing.
- 3. Add any SC formulation mix partners and agitate to ensure complete mixing.
- 4. Add BRAZEN ALL IN™ (EC) herbicide and agitate to ensure complete mixing.
- 5. Add any/additional EC formulation mix partners and agitate to ensure complete mixing.
- 6. Fill the tank to ¾ the required amount of water.
- 7. Add any solution (SN) formulation mix partners and agitate to ensure complete mixing.
- 8. Finish filling the sprayer with water, maintaining good agitation.
- 9. After any break in spraying operations, agitate thoroughly before spraying again.
- 10. Spray the herbicide suspension the same day as mixing.



CLICK HERE FOR FULL PRODUCT DETAILS.



Always read and follow label directions.



Always read and follow label directions. Toll-free: 1.855.264.6262 | Website: ADAMA.COM BADGE®, BUMPER® 432 EC, ESTEEM®, FORCEFIGHTER® M, OUTSHINE® and TOPLINE® ALL IN™, OUTSHINE ALL IN™ and TOPNOTCH™ are trademarks of ADAMA Agricultural Solutions Canada Ltd. © 2024 ADAMA Agricultural Solutions Canada Ltd.

Quick Tips

Under ideal growing conditions expect to see onset of symptomology in 3-5 days.

¹Always consult the label of the broadleaf herbicide prior to use.

² For control of common ragweed and suppression of round-leaved mallow.